

(pegloticase)

KRYSTEXXA Infusion Orders



Patient Name _____ DOB _____ Gender: M F

Phone _____

DIAGNOSIS: *Please provide IDC-10 Code*

- _____ Chronic Gout
- _____ (Other)

PRE-MEDICATION:

- | | |
|---|---|
| <input type="checkbox"/> Tylenol 1000mg PO | <input type="checkbox"/> Diphenhydramine 25mg IVP |
| <input type="checkbox"/> Cetirizine 10mg PO | <input type="checkbox"/> Solu-Cortef 100mg IVP |
| <input type="checkbox"/> _____ Other | <input type="checkbox"/> _____ Other |

KRYSTEXXA ORDERS:

DOSAGE/FREQUENCY:

- 8mg IV every 2 weeks
- Emergency Orders and Treatment per Specialty Infusion Center Anaphylaxis Policy*

PATIENT WEIGHT

_____ LBS
_____ KG

PRE-MEDICATION PER PRESCRIBING INFORMATION :

- Solu-medrol 125mg IV
- Diphenhydramine 25mg PO

NOTES:

Nursing Orders

- Assess patient prior to each infusion
- Provide patient/caregiver education related to disease process, therapy, infection control, drug use, side effects and precautions, emergency plan
- Weigh patient every visit and record
- Monitor vital signs initially and throughout infusion per manufacturer recommendations
- Establish and maintain IV access / may use central venous access if appropriate and maintain per Specialty Infusion Center policy
- Flush IV with 3ml-5ml of saline before and after every IV medication and PRN
- Administer infusion medication/injection as ordered
- If mild itching, slow down the infusion and monitor patient closely
- If anaphylactic reaction, stop infusion, implement emergency medications, and plan and call 911
- When infusion complete, flush IV with 3ml to 5ml saline flush or 50ml bag of saline and discontinue IV per INS Standards
- Evaluate effectiveness of infused therapy on disease process
- Lab work per physician when ordered

ORDERING PROVIDER:

Signature _____ Printed Name: _____ Date _____

Practice _____ Phone _____ Fax _____

*****PLEASE FAX COMPLETED ORDER TO SPECIALTY INFUSION CENTER 231-600-7058*****